

BRITISH GEOLOGICAL SURVEY

Lerwick Observatory

Monthly Magnetic Bulletin

February 2009

09/02/LE



SHETLAND ISLANDS



British Geological Survey

NATURAL ENVIRONMENT RESEARCH COUNCIL

LERWICK OBSERVATORY MAGNETIC DATA

1. Introduction

This bulletin is published to meet the needs of both commercial and academic users of geomagnetic data. Magnetic observatory data is presented as a series of plots of one-minute, hourly and daily values, followed by tabulations of monthly values, geomagnetic activity indices and reports of rapid variations. The operation of the observatory and presentation of data are described in the rest of this section.

Enquiries about the data should be addressed to:

National Geomagnetic Service
British Geological Survey
Murchison House, West Mains Road
Edinburgh EH9 3LA
Scotland, UK

Tel: +44 (0) 131 667 1000
Fax: +44 (0) 131 650 0265
E-mail: orba@bgs.ac.uk
Internet: www.geomag.bgs.ac.uk

2. Position

Lerwick Observatory, one of the three geomagnetic observatories operated and maintained in the UK by BGS, is situated on a ridge of high ground about 2.5 km to the SW of the port of Lerwick in Shetland. The observatory co-ordinates are:

Geographic: 60°07.98'N 358°48.96'E
Geomagnetic: 61°59.52'N 88°55.8'E
Height above mean sea level: 85 m

The geomagnetic co-ordinates are calculated using the 10th generation International Geomagnetic Reference Field at epoch 2009.5.

3. The Observatory Operation

3.1 GDAS

The observatory operates under the control of the Geomagnetic Data Acquisition System (GDAS), which was developed by BGS staff, installed in 2002, and became fully operational in January 2003. The system operates under the control of data acquisition software running on QNX computers, which control the data logging and communications.

There are two sets of sensors used for making magnetic measurements. A tri-axial linear-core fluxgate magnetometer, manufactured by the Danish Meteorological Institute, is used to measure the variations in the horizontal (H) and vertical (Z) components of the field. The third sensor is oriented perpendicular to these, and measures variations, which are proportional to the changes in declination (D). Measurements are made at a rate of 1 Hz.

In addition to the fluxgate sensors there is a proton precession magnetometer (PPM) making measurements of the absolute total field intensity (F) at a rate of 0.1Hz.

The raw unfiltered data are retrieved automatically via Internet connections to the BGS office in Edinburgh in near real-time. The fluxgate data are filtered to produce one-minute values using a 61-point cosine filter whilst the total field intensity samples are filtered using a 7-point cosine filter. These one-minute values are used to update the Geomagnetism Information and Forecast Service (GIFS), an on-line information system accessed via the World Wide Web at the address given in Section 1.1. GIFS also provides information on geomagnetic and solar activity.

3.2 Back-up Systems

There are two other fully independent identical systems, GDAS 2 and GDAS 3, operating at the observatory. The data from these are also processed in near real-time and used for quality control purposes. They can also be used to fill any gaps or replace any corrupt values in the primary system, GDAS 1.

3.3 Absolute Observations

The GDAS fluxgate magnetometers accurately measure variations in the components of the geomagnetic field, but not the absolute magnitudes. Two sets of absolute measurements of the field are made manually once per week. A fluxgate sensor mounted on a theodolite is used to determine D and inclination (I); the GDAS PPM measurements, with a site difference correction applied, are used for F . The absolute observations are used in conjunction with the GDAS variometer measurements to produce a continuous record of the absolute values of the geomagnetic field elements as if they had been measured at the observatory reference pillar.

4. Observatory Results

The data presented in the bulletin are in the form of plots and tabulations described in the following sections.

4.1 Absolute Observations

The absolute observation measurements made during the month are tabulated. Also included are the corresponding baseline values, which are the differences between the absolute measurements and the variometer measurements of D , H and Z (in the sense absolute–variometer). These are also plotted (markers) along with the derived preliminary daily baseline values (line) throughout the year. Daily mean differences between the measured absolute F and the F computed from the baseline corrected H and Z values are plotted in the fourth panel (in the sense measured–derived). The bottom panel shows the daily mean temperature in the fluxgate chamber.

4.2 Summary magnetograms

Small-scale magnetograms are plotted which allow the month's data to be viewed at a glance. They are plotted 16 days a page and show the variations in D , H and Z . The scales are shown on the right-hand side of the page. On disturbed days the scales are multiplied by a factor, which is indicated above the panel for that day. The variations are centred on the monthly mean value, shown on the left side of the page.

4.3 Magnetograms

The daily magnetograms are plotted using one-minute values of D , H and Z from the fluxgate sensors, with any gaps filled using back-up data. The magnetograms are plotted to a variable scale; scale bars are shown to the right of each plot. The absolute level (the monthly mean value) is indicated on the left side of the plots.

4.4 Hourly Mean Value Plots

Hourly mean values of D , H and Z for the past 12 months are plotted in 27-day segments corresponding to the Bartels solar rotation number. Magnetic disturbances associated with active regions on the surface of the Sun may recur after 27 days: the same is true for geomagnetically quiet intervals. Plotting the data in this way highlights this recurrence, and also illustrates seasonal and diurnal variations throughout the year.

4.5 Daily and Monthly Mean Values

Daily mean values of D , H , Z and F are plotted throughout the year. In addition, a table of monthly mean values of all the geomagnetic elements is provided. These values depend on accurate specification of the fluxgate sensor baselines. Provisional and definitive values are indicated in the table as **P** or **D** respectively. It is anticipated that provisional values will not be altered by more than a few nT or tenths of arcminutes before being made definitive.

4.6 Geomagnetic activity indices

The Observatory K index. This summarises geomagnetic activity at an observatory by assigning a code, an integer in the range 0 to 9, to each 3-hour Universal Time (UT) interval. The index for each 3-hour UT interval is determined from the ranges in H and in D (scaled in nT), with allowance made for the regular (undisturbed) diurnal variation. The conversion from range to an index value is made using a quasi-logarithmic scale, with the scale values dependent on the geomagnetic latitude of the observatory. The K index retains the local time (LT) and seasonal dependence of activity associated with the position of the observatory.

The provisional aa index. A number of 3-hour geomagnetic indices are computed by combining K indices from networks of observatories to characterise global activity levels and to eliminate LT and seasonal effects. The simplest of these is the aa index, computed using the K indices from two approximately antipodal observatories: Hartland in the UK and Canberra in Australia. The aa index is calculated from linearisations of the Hartland and Canberra K indices, and has units of nT. The daily mean value of aa (denoted Aa), the mean values of aa for the intervals 00-12UT and 12-24UT and the daily mean values for Hartland alone (Aa_n) and Canberra alone (Aa_s) are tabulated.

Although the aa index is based on data from only two observatories, provided averages over 12 hours or longer are used, the index is strongly correlated with the ap and am indices, which are derived using data from more extensive observatory networks.

The aa indices listed in this publication are provisional only; the definitive values are published by the International Service for Geomagnetic Indices, CRPE/CNET - CNRS, 4 Avenue de Neptune, F-94107 Saint Maur Cedex, France.

4.7 Rapid Variations

Charged particles stream from the Sun in the solar wind. The solar wind interacts with the


geomagnetic field to create a cavity, the magnetosphere, in which the field is confined. When a region of enhanced velocity and/or density in the solar wind arrives at the dayside boundary of the magnetosphere (at about 10 earth radii) the boundary is pushed towards the Earth. Currents set up on the boundary of the magnetosphere can cause an abrupt change in the geomagnetic field measured on the ground and this is recorded on observatory magnetograms as a Sudden Impulse (SI). If, following an SI, there is a change in the rhythm of activity, the SI is termed a Storm Sudden Commencement (SSC). A classical magnetic storm exhibiting initial, main and recovery phases (shown by, for instance, the *Dst* ring current index) can often occur after a SSC, in which case the start of the storm is taken as the time of the SSC.

Solar flares, seen at optical wavelengths as a sudden brightening of a small region of the Sun's surface, are also responsible for increased X-ray emissions. The X-rays cause increased ionisation in the ionosphere, which leads to absorption of short-wave radio signals. On an observatory magnetogram a Solar Flare Effect (SFE), or "crochet" may be observed. This is an enhancement to the diurnal variation of the order of 10 nT, lasting about an hour.

**This product includes mapping data licensed from Ordnance Survey with the permission of HMSO © Crown copyright. All rights reserved.
Licence Number: 100017897/2009**

© NERC 2009


Lerwick 2009




Lerwick

February

2009





Lerwick


February

2009

Date: 01-02-2009


Lerwick

Day number: 032



Date: 02-02-2009


Day number: 033



Date: 03-02-2009


Lerwick

Day number: 034



Date: 04-02-2009


Day number: 035



Date: 05-02-2009


Lerwick

Day number: 036



Date: 06-02-2009


Day number: 037



Date: 07-02-2009


Lerwick

Day number: 038



Date: 08-02-2009


Day number: 039



Date: 09-02-2009


Lerwick

Day number: 040



Date: 10-02-2009


Day number: 041



Date: 11-02-2009


Lerwick

Day number: 042



Date: 12-02-2009


Day number: 043



Date: 13-02-2009


Lerwick

Day number: 044



Date: 14-02-2009


Day number: 045



Date: 15-02-2009


Lerwick

Day number: 046



Date: 16-02-2009


Day number: 047



Date: 17-02-2009


Lerwick

Day number: 048



Date: 18-02-2009


Day number: 049



Date: 19-02-2009


Lerwick

Day number: 050



Date: 20-02-2009


Day number: 051



Date: 21-02-2009


Lerwick

Day number: 052



Date: 22-02-2009


Day number: 053



Date: 23-02-2009


Lerwick

Day number: 054



Date: 24-02-2009


Day number: 055



Date: 25-02-2009


Lerwick

Day number: 056



Date: 26-02-2009


Day number: 057



Date: 27-02-2009


Lerwick

Day number: 058




Date: 28-02-2009

Day number: 059




Lerwick Observatory: Declination (degrees)




Hourly Mean Values Plotted By Bartels Solar Rotation Number

Lerwick Observatory: Horizontal Intensity (nT)




Hourly Mean Values Plotted By Bartels Solar Rotation Number

Lerwick Observatory: Vertical Intensity (nT)



Lerwick Observatory 2009



Monthly Mean Values for Lerwick Observatory 2009

| Month | <i>D</i> | <i>H</i> | <i>I</i> | <i>X</i> | <i>Y</i> | <i>Z</i> | <i>F</i> |
|----------|-----------|----------|-----------|----------|----------|----------|----------|
| January | -3° 14.2′ | 14997 nT | 72° 48.4′ | 14973 nT | -847 nT | 48470 nT | 50738 nT |
| February | -3° 13.3′ | 14999 nT | 72° 48.4′ | 14975 nT | -843 nT | 48471 nT | 50738 nT |

Note

i. The values shown here are provisional.

LERWICK RAPID VARIATIONS

SI and SSCs

| Date | Time (UT) | Type | Quality | H (nT) | D (min) | Z (nT) |
|----------|-----------|------|---------|--------|---------|--------|
| 03-02-09 | 20 12 | SSC* | B | 9.2 | -0.56 | -3.4 |

Notes:

An asterisk (*) indicates that the principal impulse was preceded by a smaller reversed impulse.

The quality of the event is classified as follows:

A = very distinct

B = fair, ordinary, but unmistakable

C = doubtful

The amplitudes given are for the first chief movement of the event.

SFEs

| Date | Universal Time | | | H (nT) | D (min) | Z (nT) |
|------|----------------|---------|-----|--------|---------|--------|
| | Start | Maximum | End | | | |
| NONE | | | | | | |

Note:

The amplitudes given are for the first chief movement of the event.

INDICES OF GEOMAGNETIC ACTIVITY

Lerwick Observatory

February 2009

| Day | K - INDICES FOR THREE-HOUR INTERVAL | | | | | | | | SUM |
|-----|-------------------------------------|-------|-------|-------|-------|-------|-------|-------|-----|
| | 00-03 | 03-06 | 06-09 | 09-12 | 12-15 | 15-18 | 18-21 | 21-24 | |
| 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 2 |
| 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 3 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 3 | 4 |
| 4 | 3 | 1 | 1 | 2 | 2 | 3 | 3 | 2 | 17 |
| 5 | 3 | 1 | 0 | 0 | 1 | 1 | 0 | 2 | 8 |
| 6 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 |
| 7 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| 8 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 9 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 2 |
| 10 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 11 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| 12 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 2 |
| 13 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 2 |
| 14 | 1 | 1 | 2 | 3 | 3 | 3 | 1 | 4 | 18 |
| 15 | 2 | 2 | 1 | 1 | 2 | 2 | 3 | 3 | 16 |
| 16 | 1 | 0 | 0 | 0 | 1 | 2 | 0 | 1 | 5 |
| 17 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 18 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 2 | 4 |
| 19 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 1 |
| 20 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 2 | 4 |
| 21 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 |
| 22 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 1 | 3 |
| 23 | 0 | 0 | 0 | 1 | 1 | 2 | 2 | 3 | 9 |
| 24 | 3 | 2 | 1 | 1 | 0 | 0 | 0 | 0 | 7 |
| 25 | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 0 | 3 |
| 26 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 2 |
| 27 | 1 | 1 | 1 | 3 | 2 | 2 | 2 | 2 | 14 |
| 28 | 2 | 2 | 1 | 0 | 0 | 1 | 0 | 2 | 8 |
| | | | | | | | | | |

| | | | | | | | | | |
|--|----|----|----|----|-----|-----|-----|-----|------|
| Lower bound (nT) for the range for each index value at Lerwick Observatory | | | | | | | | | |
| K-Index | | | | | | | | | |
| 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| 0 | 10 | 20 | 40 | 80 | 140 | 240 | 400 | 660 | 1000 |

The aa Index

| Date | Day | K-North | K-South | (a) | (b) | (c) | (d) | (e) |
|----------------------|-----|-----------------|-----------------|-----|-----|-----|-----|-----|
| 01-02-09 | 32 | 1 1 1 0 1 1 0 1 | 0 1 1 1 0 0 1 0 | 6 | 5 | 6 | 5 | 6 |
| 02-02-09 | 33 | 0 0 0 0 1 1 0 0 | 1 1 0 0 0 0 0 0 | 4 | 4 | 4 | 4 | 4 |
| 03-02-09 | 34 | 0 0 0 0 1 1 2 4 | 0 0 1 0 0 0 1 3 | 12 | 8 | 3 | 17 | 10 |
| 04-02-09 | 35 | 4 1 1 3 2 3 4 2 | 3 1 2 3 4 3 3 2 | 28 | 29 | 24 | 33 | 29 |
| 05-02-09 | 36 | 3 2 1 1 1 1 1 2 | 2 2 0 2 2 2 0 1 | 13 | 12 | 14 | 10 | 12 |
| 06-02-09 | 37 | 2 0 0 0 0 0 0 1 | 0 0 0 0 0 0 0 0 | 5 | 2 | 4 | 3 | 4 |
| 07-02-09 | 38 | 0 2 0 0 1 1 0 0 | 1 1 0 0 1 1 0 0 | 5 | 5 | 6 | 5 | 5 |
| 08-02-09 | 39 | 0 0 0 0 1 1 0 1 | 0 1 1 0 0 0 0 0 | 4 | 4 | 4 | 4 | 4 |
| 09-02-09 | 40 | 1 0 0 0 1 1 0 2 | 1 1 1 0 0 1 0 0 | 6 | 5 | 5 | 6 | 6 |
| 10-02-09 | 41 | 0 0 0 0 1 1 1 0 | 1 1 1 0 1 1 0 1 | 4 | 7 | 4 | 6 | 5 |
| 11-02-09 | 42 | 2 0 0 0 1 0 1 0 | 1 1 1 0 1 1 1 1 | 5 | 7 | 6 | 6 | 6 |
| 12-02-09 | 43 | 0 1 0 1 1 2 1 0 | 2 2 0 2 2 2 0 0 | 7 | 11 | 9 | 9 | 9 |
| 13-02-09 | 44 | 0 0 0 0 1 1 1 1 | 1 0 0 0 1 1 1 1 | 5 | 6 | 3 | 8 | 5 |
| 14-02-09 | 45 | 1 2 3 3 4 4 1 4 | 2 3 4 4 5 2 1 2 | 34 | 39 | 32 | 41 | 36 |
| 15-02-09 | 46 | 2 3 1 2 3 2 3 3 | 1 3 4 3 4 2 1 1 | 23 | 28 | 25 | 25 | 25 |
| 16-02-09 | 47 | 2 0 0 0 1 2 1 1 | 1 1 1 1 1 2 1 1 | 8 | 9 | 7 | 10 | 8 |
| 17-02-09 | 48 | 0 0 0 0 0 1 0 1 | 0 0 0 0 0 0 0 1 | 4 | 3 | 2 | 4 | 3 |
| 18-02-09 | 49 | 2 0 1 0 1 1 2 3 | 1 0 0 0 0 1 0 1 | 11 | 5 | 5 | 10 | 8 |
| 19-02-09 | 50 | 0 0 0 0 0 1 0 1 | 0 0 0 1 0 2 0 1 | 4 | 6 | 3 | 6 | 5 |
| 20-02-09 | 51 | 1 1 1 0 1 1 3 3 | 1 1 1 1 1 2 1 1 | 13 | 9 | 7 | 15 | 11 |
| 21-02-09 | 52 | 1 0 0 1 1 1 1 1 | 0 0 1 2 1 1 0 2 | 6 | 8 | 6 | 8 | 7 |
| 22-02-09 | 53 | 0 1 1 1 2 1 1 2 | 1 1 1 2 2 0 0 2 | 9 | 10 | 8 | 11 | 9 |
| 23-02-09 | 54 | 1 0 0 1 2 3 2 3 | 1 1 0 1 3 3 1 1 | 14 | 13 | 6 | 22 | 14 |
| 24-02-09 | 55 | 3 3 2 1 1 1 1 1 | 1 2 1 1 2 1 0 1 | 15 | 9 | 16 | 8 | 12 |
| 25-02-09 | 56 | 2 1 1 1 1 1 2 1 | 1 0 0 1 0 0 0 0 | 10 | 4 | 7 | 6 | 7 |
| 26-02-09 | 57 | 1 1 0 1 1 1 1 1 | 1 1 0 0 1 1 0 1 | 7 | 6 | 6 | 7 | 6 |
| 27-02-09 | 58 | 2 1 2 3 3 3 2 3 | 2 2 1 4 3 2 2 1 | 23 | 22 | 21 | 23 | 22 |
| 28-02-09 | 59 | 3 2 2 0 1 1 2 2 | 1 2 1 0 0 0 1 0 | 14 | 6 | 13 | 8 | 10 |
| Monthly mean value = | | | 10.4 | | | | | |

- (a) The northern daily mean value, Aa_n
- (b) The southern daily mean value, Aa_s
- (c) The mean value of aa for the interval 00-12 UT
- (d) The mean value of aa for the interval 12-24 UT
- (e) The daily mean value of aa (Aa)

Notes

- i. The values are rounded to the nearest integer.
- ii. The units of the aa index are nT.
- iii. The values shown here are provisional. The definitive values are computed and published by the International Service for Geomagnetic Indices, Paris